


This article was downloaded by: [Qingdao University]

On: 09 October 2014, At: 19:36

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Applicable Analysis: An International Journal

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gapa20>

Spaces of harmonic functions with boundary values in

Qixiang Yang^a, Tao Qian^b & Pengtao Li^c

^a School of Mathematics and Statistics, Wuhan University, Wuhan, 430072China.

^b Faculty of Science and Technology, Department of Mathematics, University of Macau, Macau, China.

^c College of Mathematics, Qingdao University, Qingdao, Shandong, 266071China.

Published online: 07 Oct 2014.

To cite this article: Qixiang Yang, Tao Qian & Pengtao Li (2014): Spaces of harmonic functions with boundary values in , *Applicable Analysis: An International Journal*, DOI: [10.1080/00036811.2014.959441](https://doi.org/10.1080/00036811.2014.959441)

To link to this article: <http://dx.doi.org/10.1080/00036811.2014.959441>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Spaces of harmonic functions with boundary values in $Q_{p,q}^\alpha$

Qixiang Yang^a, Tao Qian^b and Pengtao Li^{c*}

^aSchool of Mathematics and Statistics, Wuhan University, Wuhan 430072, China; ^bFaculty of Science and Technology, Department of Mathematics, University of Macau, Macau, China; ^cCollege of Mathematics, Qingdao University, Qingdao, Shandong 266071, China

Communicated by Y. Xu

(Received 25 August 2014; accepted 26 August 2014)

In this paper, we apply wavelets to study two classes of function spaces of harmonic functions: the weighted Besov spaces $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ and Carleson spaces $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$. By a reproducing formula, we prove that the elements in these harmonic function spaces can be characterized by the Poisson integral of the functions in the Besov-Q spaces $Q_{p,q}^\alpha(\mathbb{R}^n)$.

Keywords: wavelet; weighted Besov spaces; Carleson measures; Besov-Morrey space

AMS Subject Classifications: 42B35; 42C40

1. Introduction

In this paper, we use wavelets to study the spaces of harmonic functions with boundary values in Besov-Q spaces $Q_{p,q}^\alpha(\mathbb{R}^n)$. It is well known that for a measurable function on \mathbb{R}^n , the Poisson integral $P_t f$ gives a harmonic extension of f . In the literature, Poisson integral is used to describe the relation between the harmonic function spaces on \mathbb{R}_+^{n+1} and their boundary values. Fabes et al. [1] characterized the spaces $HMO(\mathbb{R}_+^{n+1})$ with trace in $BMO(\mathbb{R}^n)$. Precisely, they proved the following result:

$$u \in HMO(\mathbb{R}_+^{n+1}) \iff u = P_t * f \text{ for some } f \in BMO(\mathbb{R}^n). \quad (1.1)$$

See [1, Theorem 1.0].

By wavelet methods, we will establish the following relations among the Besov-Q spaces $Q_{p,q}^\alpha(\mathbb{R}^n)$, the wavelet spaces $W_{p,q}^\alpha(\mathbb{R}^n)$, the weighted Besov spaces $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ and the Carleson spaces $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$:

$$Q_{p,q}^\alpha \xrightarrow{\text{Theorem 2.8}} W_{p,q}^\alpha \ni f(x) \xleftrightarrow[\text{Theorem 4.2}]{\text{Theorem 4.4}} f(x, t) \in C_{p,q}^\alpha \xrightarrow{\text{Theorem 3.4}} H_{p,q}^{\alpha,\lambda}. \quad (1.2)$$

*Corresponding author. Email: ptli@qdu.edu.cn

We give the definitions of $Q_{p,q}^\alpha(\mathbb{R}^n)$ and $W_{p,q}^\alpha(\mathbb{R}^n)$ in Section 2. The definitions of the harmonic function spaces $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ and $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$ can be found in Section 3. Precisely, in this paper, we show the following results.

THEOREM 1.1 *Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. The following five statements are equivalent:*

- (i) $f(x) \in Q_{p,q}^\alpha(\mathbb{R}^n)$.
- (ii) $f(x) \in W_{p,q}^\alpha(\mathbb{R}^n)$.
- (iii) $P_t f(x) \in H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$, $\exists \lambda > n(1 - \frac{q}{p})$.
- (iv) $P_t f(x) \in H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$, $\forall \lambda > n(1 - \frac{q}{p})$.
- (v) $P_t f(x) \in C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$.

The significance of these spaces is that for particular choices of the parameters p, q and α , one obtains various classical function spaces, such as the Bergman spaces, the Bloch spaces, the Besov spaces, the BMO spaces and the Q spaces. We give the following space structure table to clarify the relation between these spaces and $Q_{p,q}^\alpha(\mathbb{R}^n)$:

$\alpha \in [0, 1), 1 \leq p = q \leq \infty,$	Besov spaces [2]
$\alpha = 0, p = \infty, q = 2,$	BMO space [3]
$\alpha \in (0, \min(1, \frac{n}{2})), p = 2n/\alpha, q = 2,$	Q-spaces Q_α [4]
$\alpha = 0, p > 2, q = 2,$	Morrey spaces $L^{2,\lambda}$ [5]
$\alpha = 0, p = q = 1,$	real Bergman space [6,7]
$\alpha = 0, p = q = \infty,$	real Bloch space [6]

In the proof of Theorem 1.1, we need to overcome two difficulties:

On one hand, for a harmonic function $F(x, t)$, its boundary value may not be obtained via pointwise limits. In the paper, we use an alternative way to define the boundary value of functions in $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$. By a reproducing formula (2.3), we define the boundary value of $f(x, t)$ via (2.4).

On the other hand, to characterize $Q_{p,2}^\alpha(\mathbb{R}^n)$ by the Poisson kernel and the heat semi-groups, one of the main methodologies is the Fourier transform. See [8]. However for the spaces $Q_{p,q}^\alpha$ with $q \neq 2$, Fourier transform does not work. For functions $f \in Q_{p,q}^\alpha$, to surmount this obstacle, we use regular wavelets to estimate the Poisson kernel $P_t(x)$.

Now we give an outline of the proof of Theorem 1.1.

- (1) The equivalence (i) and (ii) is well known. We list it as Theorem 2.8. See Section 2.4 and the references.[9–11]
- (2) In Section 3, we prove that $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1}) = C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$. In fact, our result implies that $f(x, t) \in H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ if and only if

$$d\mu =: |\nabla f(x, t)|^q t^{q-1-q\alpha} dx dt$$

is a $(1 - q/p)$ -Carleson measure. See Theorem 3.4 for the equivalence of (iii), (iv) and (v).

- (3) Let P_t be the Poisson kernel. In Lemma 2.5, we estimate the wavelet coefficients of the function $\frac{\partial}{\partial x_i} P_t(x - y)$. With the help of Lemma 2.5 and Theorem 2.8, we can get the following inclusion relation in Section 4.1:

$$P_t * (Q_{p,q}^\alpha(\mathbb{R}^n)) \subseteq C_{p,q}^\alpha(\mathbb{R}_+^{n+1}).$$

This gives (ii) \Rightarrow (v). In Section 4.2, we prove that $f(x, t) \in C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$ can be represented as the Poisson integral $P_t * f(x)$, where f is an element in $W_{p,q}^\alpha(\mathbb{R}^n)$. See Theorem 4.2 for a proof of (v) \Rightarrow (ii).

Remark: In a recent paper, by a different method, Wang-Xiao [24] obtain a extension of Campanato-Sobolev spaces $Q_{\lambda,2}^s$ via the fractional heat semigroups. We also refer the reader to Jiang-Xiao-Yang [25] for further information on this topic.

Some notations:

- $U \approx V$ represents that there is a constant $c > 0$ such that $c^{-1}V \leq U \leq cV$ whose right inequality is also written as $U \lesssim V$. Similarly, one writes $V \gtrsim U$ for $V \geq cU$.
- For convenience, the positive constants C may change from one line to another and usually depend on the dimension n , α , β and other fixed parameters. The Schwartz class of rapidly decreasing functions and its dual will be denoted by $\mathcal{S}(\mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{R}^n)$, respectively. For $f \in \mathcal{S}(\mathbb{R}^n)$, \widehat{f} means the Fourier transform of f .

2. Preliminaries

2.1. Regular Daubechies wavelets

We present some preliminaries on Daubechies' wavelets Φ^ϵ , $\epsilon = 0$ or 1 , and refer the reader to [6,12] and [13] for further information. Let

$$\begin{cases} E_n = \{0, 1\}^n \setminus \{0\}; \\ F_n = \{(\epsilon, k) : \epsilon \in E_n, k \in \mathbb{Z}^n\}; \\ \Lambda_n = \{(\epsilon, j, k), \epsilon \in E_n, j \in \mathbb{Z}, k \in \mathbb{Z}^n\}, \end{cases}$$

We will use the real-valued regular Daubechies' wavelets. Let C^m denote the smooth function spaces with all the derivatives up to the order m , and being bounded. In this paper, we assume there exist two sufficiently large integers m and M such that

- (i) For any $\epsilon \in E_n$, $\text{supp}\Phi^\epsilon \subset [-2^M, 2^M]^n$;
- (ii) $\Phi^\epsilon \in C^m([-2^M, 2^M]^n)$;
- (iii) For $|\alpha| \leq m$, $\int x^\alpha \Phi^\epsilon(x) dx = 0$.

For $(\epsilon, j, k) \in \Lambda_n$, let

$$\Phi_{j,k}^\epsilon(x) = 2^{jn/2} \Phi^\epsilon(2^j x - k).$$

The set $\{\Phi_{j,k}^\epsilon, (\epsilon, j, k) \in \Lambda_n\}$ forms a wavelet basis. For any $\epsilon \in \{0, 1\}^n$, $k \in \mathbb{Z}^n$ and a function f on \mathbb{R}^n , we write $f_{j,k}^\epsilon = \langle f, \Phi_{j,k}^\epsilon \rangle$. The following result is well known.

LEMMA 2.1 *The Daubechies wavelets $\{\Phi_{j,k}^\epsilon\}_{(\epsilon,j,k)\in\Lambda_n}$ form an orthogonal basis of $L^2(\mathbb{R}^n)$. Consequently, for any $f \in L^2(\mathbb{R}^n)$, the following wavelet decomposition holds in the L^2 convergence sense:*

$$f = \sum_{(\epsilon,j,k)\in\Lambda_n} f_{j,k}^\epsilon \Phi_{j,k}^\epsilon.$$

2.2. Poisson extension and boundary value

We first construct some functions with special compact supports in order to define boundary limits of harmonic functions.

LEMMA 2.2 *Fix $m \in \mathbb{N}$. There exist a constant $C_0 > 0$ and two radial real-valued functions $\phi \in C^{2m+8}(B(0, 1))$ and $\Phi \in C^{4m+8}(B(0, 1))$ such that*

- (i) $\phi(x) = (-\Delta)^m \Phi(x)$;
- (ii) $\int_0^\infty (\widehat{\phi}(t\xi))^2 \frac{dt}{t} = 1, \forall \xi \neq 0$;
- (iii) $C_0 \int_0^\infty \widehat{\phi}(t) e^{-t} \frac{dt}{t} = 1$.

Proof It is easy to choose a radial real-valued function $\Psi \in C^{4m+8}(B(0, 1))$ such that

$$\int_0^\infty t^{2m} \widehat{\Psi}(t) e^{-t} \frac{dt}{t} \neq 0.$$

Let

$$C_\Psi = \int_0^\infty t^{4m} |\xi|^{4m} |\widehat{\Psi}(t\xi)|^2 \frac{dt}{t}$$

and

$$\Phi(x) = (C_\Psi)^{-\frac{1}{2}} \Psi(x).$$

□

Let $C_n = \Gamma((n+1)/2)/\pi^{(n+1)/2}$ and let

$$\begin{cases} P(x) = \frac{C_n}{(1+|x|^2)^{(n+1)/2}}; \\ P_t(x) = t^{-n} P\left(\frac{x}{t}\right) = \frac{C_n t}{(t^2+|x|^2)^{(n+1)/2}}. \end{cases}$$

Let f be any measurable function on \mathbb{R}^n satisfying

$$\int_{\mathbb{R}^n} \frac{|f(x)|}{1+|x|^{n+1}} dx < \infty. \quad (2.1)$$

The Poisson integral of f is defined by

$$f(x, t) = \int_{\mathbb{R}^n} P_t(x-y) f(y) dy.$$

Let $A(\mathbb{R}^n) = \{f(x) : (1+|x|)^{n+1} \partial_x^\alpha f \in L^\infty, \forall \alpha \in \mathbb{N}^n\}$ and denote $A'(\mathbb{R}^n)$ be the dual space of $A(\mathbb{R}^n)$. For any function $g \in \mathcal{S}(\mathbb{R}^n)$, we know that $P_t g(x) \in A(\mathbb{R}^n) \forall t \geq 0$.

Hence, for any distribution $f \in A'(\mathbb{R}^n)$, f can be extended formally to a harmonic function $P_t f(x)$ as

$$f(x, t) = e^{-t(-\Delta)^{\frac{1}{2}}} f(x) = P_t * f(x), \tag{2.2}$$

For any $t \geq 0$, $f(x, t)$ is a distribution. That is to say, if f does not satisfy (2.1), we can still define the Poisson extension $P_t f$.

Example 2.3 Let δ be the Dirac function. It is well known that δ is not measurable. However, it is obvious that

$$P_t \delta(x) = P_t(x).$$

For the harmonic function $f(x, t) =: P_t(x)$, we have

$$\lim_{t \rightarrow 0} f(x, t) = 0, \quad \forall x \neq 0.$$

But we know, as $t \rightarrow 0$, $P_t(x)$ converges to $\delta(x)$ in the sense of distribution.

Example 2.3 implies that for a general harmonic function $f(x, t)$, its boundary value may not be defined in the sense of the pointwise limit as $t \rightarrow 0$. In Lemma 2.2, we use some compactly supported function to pull back the harmonic functions to some boundary functions.

Let ϕ be the function obtained in Lemma 2.2. Write $\phi_t(x) = t^{-n} \phi(\frac{x}{t})$ with $\widehat{\phi}_t(\xi) = \widehat{\phi}(t\xi)$. From (iii) of Lemma 2.2, we can deduce that

$$\widehat{f}(\xi) = C_0 \int_0^\infty \widehat{\phi}(t) e^{-t} \frac{dt}{t} \widehat{f}(\xi) = C_0 \int_0^\infty \widehat{\phi}(t\xi) e^{-t|\xi|} \widehat{f}(\xi) \frac{dt}{t}. \tag{2.3}$$

By the inverse Fourier transform, we can get the following result.

PROPOSITION 2.4 *If $f \in \mathcal{S}(\mathbb{R}^n)$, then the following two identities hold point by point.*

$$f(x) = \lim_{t \rightarrow 0} P_t f(x) = C_0 \int_0^\infty \int_{\mathbb{R}^n} P_t f(y) \phi_t(x - y) \frac{dt}{t} dy.$$

By Proposition 2.4, harmonic function $f(x, t)$ can be pulled back to the trace function $f(x)$ in the sense of distribution

$$f(x) = C_0 \int_0^\infty \int_{\mathbb{R}^n} f(x - y, t) \phi_t(y) \frac{dt}{t} dy.$$

2.3. Wavelet estimates on the Poisson kernel

Let

$$P_i(x) = \frac{-(n+1)C_n x_i}{(1+|x|^2)^{(n+3)/2}}.$$

For $\epsilon = (\epsilon_1, \dots, \epsilon_n) \in \{0, 1\}^n \setminus \{0\}$, let τ_ϵ be the smallest index s such that $\epsilon_s \neq 0$. Let $P_{i,\epsilon}(x) = \partial_{x_{\tau_\epsilon}} P_i(x)$. We can see that

$$P_{i,\epsilon}(x) = \begin{cases} \frac{-(n+1)C_n(1+|x|^2-(n+3)x_i^2)}{(1+|x|^2)^{(n+5)/2}}, & i = \tau_\epsilon; \\ \frac{(n+1)(n+3)C_n x_i x_{\tau_\epsilon}}{(1+|x|^2)^{(n+5)/2}}, & i \neq \tau_\epsilon. \end{cases}$$

Let $\Phi^{\epsilon,i}(x) = \frac{\partial \Phi^\epsilon(x)}{\partial x_i}$ and

$$I_\epsilon \Phi^\epsilon(x) = \int_{-\infty}^{x_{\tau_\epsilon}} \Phi^\epsilon(x_1, \dots, x_{-1+\tau_\epsilon}, y, x_{1+\tau_\epsilon}, \dots, x_n) dy.$$

For $i = 1, 2, \dots, n$, let

$$P_{i,t}(x) = -(n+1)C_n \frac{tx_i}{(t^2 + |x|^2)^{(n+3)/2}}.$$

For $i = 1, \dots, n$ and $(\epsilon, j, k) \in \Lambda_n$, define

$$\begin{aligned} I(i, t, x, \epsilon, j, k) &= \frac{\partial}{\partial x_i} \int P_t(x-y) \Phi_{j,k}^\epsilon(y) dy \\ &=: \int_{\mathbb{R}^n} P_{i,t}(x-y) \Phi_{j,k}^\epsilon(y) dy. \end{aligned} \quad (2.4)$$

We estimate $I(i, t, x, \epsilon, j, k)$ by wavelets.

LEMMA 2.5

(i) If $2^j t > 1$, then

$$|I(i, t, x, \epsilon, j, k)| \lesssim \frac{2^{(\frac{n}{2}+2)j} t}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}}. \quad (2.5)$$

(ii) If $2^j t \leq 1$, then

$$|I(i, t, x, \epsilon, j, k)| \lesssim \begin{cases} \frac{2^{(\frac{n}{2}+2)j} t}{(1+|2^j x - k|^2)^{\frac{n+1}{2}}}, & |2^j x - k| \geq C_\Phi; \\ 2^{(\frac{n}{2}+1)j}, & |2^j x - k| < C_\Phi. \end{cases} \quad (2.6)$$

Proof

(i) For $2^j t > 1$, by the change of variable, we have

$$\begin{aligned} |I(i, t, x, \epsilon, j, k)| &\lesssim t^{-2} 2^{-j} \int_{\mathbb{R}^n} |P_{i,\epsilon,t}(x-y)| |(I_\epsilon \Phi^\epsilon)_{j,k}(y)| dy \\ &\lesssim t 2^{2j+nj/2} \int_{\mathbb{R}^n} \frac{1}{(4j t^2 + |2^j x - k - u|^2)^{\frac{n+3}{2}}} |\Phi^\epsilon(u)| du. \end{aligned}$$

Because $\text{supp } \Phi^\epsilon \subset B(0, 2^M)$, we have $|2^j x - k - u| \lesssim |2^j x - k| + 1$. On the other hand, by $2^j t > 1$, we can see that there exists a constant C large enough such that

$$C[4^j t^2 + |2^j x - k - u|^2] \geq C4^j t^2 + |2^j x - k|^2 - 2|u|^2 \geq 4^j t + |2^j x - k|^2.$$

Then, we obtain

$$|I(i, t, x, \epsilon, j, k)| \lesssim t 2^{2j+nj/2} \frac{1}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}}.$$

Now we prove (ii). If $2^j t \leq 1$, applying integration by parts, we can get

$$I(i, t, x, \epsilon, j, k) = \int_{\mathbb{R}^n} P_t(x - y) 2^j \Phi_{j,k}^{\epsilon,i}(y) dy$$

Hence, we have

$$\begin{aligned} |I(i, t, x, \epsilon, j, k)| &\lesssim \int_{\mathbb{R}^n} P_t(x - y) 2^j |\Phi_{j,k}^{\epsilon,i}(y)| dy \\ &\lesssim \int_{\text{supp } \Phi^\epsilon} \frac{2^{(2+\frac{n}{2})j} t}{(4jt^2 + |2^j x - k - u|^2)^{\frac{n+1}{2}}} |\Phi^{\epsilon,i}(u)| du. \end{aligned}$$

We distinguish two cases. If $|2^j x - k| \geq C_\Phi$, we can get

$$|2^j x - k - u| \geq |2^j x - k| - |u| \geq \frac{1}{2} |2^j x - k| \geq C_\Phi/2.$$

On the other hand, by $2^j t \leq 1$, $4jt^2 \lesssim |2^j x - k|^2$. The above estimates imply that

$$|I(i, t, x, \epsilon, j, k)| \lesssim \frac{2^{(2+\frac{n}{2})j} t}{(|2^j x - k|^2 + C_\Phi^2)^{\frac{n+1}{2}}} \lesssim \frac{2^{(\frac{n}{2}+2)j} t}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}}$$

If $|2^j x - k| \leq C_\Phi$, because $|\Phi^{\epsilon,i}(2^j y - k)| \leq C$, a direct computation gives

$$\begin{aligned} |I(i, t, x, \epsilon, j, k)| &\lesssim 2^{j(1+\frac{n}{2})} \int_{\mathbb{R}^n} \frac{t}{(t^2 + |x - y|^2)^{\frac{n+1}{2}}} |\Phi^{\epsilon,i}(2^j y - k)| dy \\ &\lesssim 2^{j(1+\frac{n}{2})}. \end{aligned}$$

This completes the proof of Lemma 2.5. □

2.4. Besov-Q spaces and wavelet characterization

Besov-Q spaces $Q_{p,q}^\alpha(\mathbb{R}^n)$ are studied in [11].

Definition 2.6 Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. The Besov-Q space $Q_{p,q}^\alpha(\mathbb{R}^n)$ is defined to be the set of all functions with

$$\sup_I (f, Q_{p,q}^\alpha)(I) =: \sup_I |I|^{\frac{q}{p}-1} \int_I \int_I \frac{|f(x) - f(y)|^q}{|x - y|^{n+q\alpha}} dx dy < +\infty,$$

where the supremum is taken over all cubes I with the edge length $\ell(I)$ and the edges parallel to the coordinate axes in \mathbb{R}^n .

For $\alpha \in (0, 1)$, $p = n/\alpha$, $q = 2$, $Q_{n/\alpha, 2}^\alpha(\mathbb{R}^n) = Q_\alpha(\mathbb{R}^n)$. Q spaces $Q_\alpha(\mathbb{R}^n)$ were studied extensively. For further information on $Q_\alpha(\mathbb{R}^n)$, we refer the reader to Dafni-Xiao [14,15], Essen et al. [4], Wu-Xie [5] and the reference therein. The space $Q_{p,q}^\alpha(\mathbb{R}^n)$ with $\alpha \in (0, 1)$ and $2 \leq q < p < \infty$ was introduced by Cui-Yang [9]. Yang-Yuan [11] established the Littlewood-Paley characterization of $Q_{p,q}^\alpha(\mathbb{R}^n)$ with the full indices as in Definition 2.6.

Let $\{\Phi_{j,k}^\varepsilon\}$ be a wavelet basis defined in Section 2.1. For any function f , let $\{f_{j,k}^\varepsilon\}$ be the wavelet coefficients of f . By Lemma 2.1, formally

$$f = \sum_{(\varepsilon,j,k) \in \Lambda_n} f_{j,k}^\varepsilon \Phi_{j,k}^\varepsilon.$$

We introduce a space which consists of $\{f_{j,k}^\varepsilon\}$ as follows.

Definition 2.7 Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. The space $W_{p,q}^\alpha(\mathbb{R}^n)$ is defined to be the set of all functions with the wavelet coefficients satisfying

$$\sup_I \left\{ (f, W_{p,q}^\alpha)(I) \right\}^{\frac{1}{q}} =: \sup_I \left\{ |I|^{\frac{q}{p}-1} \sum_{(\varepsilon,j,k) \in \Lambda_n: I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} |f_{j,k}^\varepsilon|^q \right\}^{1/q} < \infty,$$

where the supremum is taken over all dyadic cubes I .

For $\alpha \in (0, 1)$ and $2 \leq q \leq p < \infty$, the wavelet characterization of $Q_{p,q}^\alpha(\mathbb{R}^n)$ is obtained by Cui–Yang [9]. By different methods, Lin–Yang [10] and Yang–Yuan [11] improved the scope to $\alpha \in [0, \infty)$ and $1 \leq q \leq p \leq \infty$. See also [16] and [17].

THEOREM 2.8 Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. Then

$$Q_{p,q}^\alpha(\mathbb{R}^n) = W_{p,q}^\alpha(\mathbb{R}^n).$$

3. Weighted Besov spaces and Carleson measures

On the unit disc, Zhao [18] introduced a family of analytic functions on the open unit disk, denoted by $F(p, q, s)$. The spaces $F(p, q, s)$ cover many known function spaces of analytic functions: the Bloch space, Bergman spaces and weighted Dirichlet spaces. Such spaces have been studied heavily by different authors. In the latest decades, $F(p, q, s)$ have been studied extensively. We refer the reader to [5, 19–23] and the reference therein.

For $1 \leq q \leq p < \infty$, $0 \leq \alpha < \min(1, \frac{n}{q})$ and $\lambda > n(1 - \frac{q}{p})$, replacing the analytic functions by harmonic functions, we introduce a class of spaces of harmonic functions on \mathbb{R}_+^{n+1} . For $1 \leq q < \infty$, we define the gradient of $f(x, t)$ by

$$|\nabla f(x, t)|^q = \sum_{i=1}^n \left| \frac{\partial f(x, t)}{\partial x_i} \right|^q.$$

Definition 3.1 Let $1 \leq q \leq p < \infty$, $0 \leq \alpha < \min(1, \frac{n}{q})$ and $\lambda > n(1 - \frac{q}{p})$. The weighted Besov spaces $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ is defined as the space of all harmonic functions such that

$$\|f\|_{H_{p,q}^{\alpha,\lambda}} = \sup_{(y,u) \in \mathbb{R}_+^{n+1}} \left\{ (f, H_{p,q}^{\alpha,\lambda})(y, u) \right\}^{\frac{1}{q}} < +\infty,$$

where

$$\begin{aligned} (f, H_{p,q}^{\alpha,\lambda})(y, u) &= \int_{(x,t) \in \mathbb{R}_+^{n+1}} \frac{|\nabla f(x, t)|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt \\ &= \sum_{i=1}^n \int_{(x,t) \in \mathbb{R}_+^{n+1}} \frac{\left| \frac{\partial f(x,t)}{\partial x_i} \right|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt \\ &\equiv \sum_{i=1}^n (f, H_{p,q}^{\alpha,\lambda})_i(y, u). \end{aligned}$$

The Carleson box based on a cube I is defined by

$$S(I) = I \times (0, \ell(I)] = \left\{ (x, t) \in \mathbb{R}_+^{n+1} : x \in I, t \in (0, \ell(I)] \right\}.$$

A positive measure μ is called a p -Carleson measure on \mathbb{R}_+^{n+1} if

$$\sup_I \frac{\mu(S(I))}{|I|^p} < \infty.$$

Here, \sup_I indicates the supremum take over all cubes in \mathbb{R}^n . Note that $p = 1$ gives the classical Carleson measure.

Definition 3.2 Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. We define Carleson spaces $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$ as the space of all harmonic functions such that

$$\|f\|_{C_{p,q}^\alpha} = \sup_I \left\{ (f, C_{p,q}^\alpha)(I) \right\}^{\frac{1}{q}} < +\infty,$$

where

$$(f, C_{p,q}^\alpha)(I) = |I|^{\frac{q}{p}-1} \int_{S(I)} |\nabla f(x, t)|^q t^{q-1-q\alpha} dx dt \equiv \sum_{i=1}^n (f, C_{p,q}^\alpha)_i(I).$$

Remark 3.3

- (i) In the above definition 3.1, for $1 \leq p = q \leq \infty$, we can take $\lambda = 0$. Then the definition of $H_{q,q}^{\alpha,0}(\mathbb{R}_+^{n+1})$ coincides with that of $C_{q,q}^\alpha(\mathbb{R}_+^{n+1})$ in the definition 3.2. In particular, $H_{1,1}^{0,0}(\mathbb{R}_+^{n+1})$ is the classic definition of Bergman spaces and $H_{\infty,\infty}^{0,0}(\mathbb{R}_+^{n+1})$ is the classic definition of Bloch spaces. See Section 8 of chapter 6 in [6].
- (ii) For $\alpha = 0, p = \infty$ and $q = 2, C_{\infty,2}^0(\mathbb{R}_+^{n+1})$ becomes the space $HMO(\mathbb{R}_+^{n+1})$ introduced by Fabes et al. [1].

We characterize the space $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ by Carleson measure.

THEOREM 3.4 Let $1 \leq q \leq p < \infty, 0 \leq \alpha < \min(1, \frac{n}{q})$ and $\lambda > n - \frac{nq}{p}$.

$$C_{p,q}^\alpha(\mathbb{R}_+^{n+1}) = H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1}).$$

Proof Let I be the cube parallel to the coordinate axes with centre y and the edge length $\ell(I)$, and let $u = \ell(I)/2$. When $(x, t) \in S(I)$, we have $(x - y)^2 + (u + t)^2 \sim \ell(I)^2$. Therefore,

$$|I|^{\frac{q}{p}-1} \int_{S(I)} |\nabla f(x, t)|^q t^{q-1-q\alpha} dx dt \lesssim \int_{(x,t) \in \mathbb{R}_+^{n+1}} \frac{|\nabla f(x, t)|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt.$$

Conversely, for any fixed (y, u) , let I be the cube parallel to the coordinate axes with centre y and the edge length $2u$. For nonnegative integer m , we use I_m to denote the cubes with the same centre as I and the length $2^m \ell(I)$.

$$\begin{aligned} I_{y,u} &\equiv \int_{(x,t) \in \mathbb{R}_+^{n+1}} \frac{|\nabla f(x, t)|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt \\ &\lesssim \int_{(x,t) \in S(I)} \frac{|\nabla f(x, t)|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt \\ &\quad + \sum_{m=0}^{\infty} \int_{S(I_{m+1}) \setminus S(I_m)} \frac{|\nabla f(x, t)|^q t^{q-1-q\alpha} u^{\frac{nq}{p}-n+\lambda}}{|(x-y)^2 + (u+t)^2|^{\frac{\lambda}{2}}} dx dt. \end{aligned}$$

When $(x, t) \in S(I)$, we have $(x - y)^2 + (u + t)^2 \sim \ell(I)^2$. If $(x, t) \in S(I_{m+1}) \setminus S(I_m)$, then $(x - y)^2 + (u + t)^2 \sim 2^{2m} \ell(I)^2$. Therefore,

$$\begin{aligned} I_{y,u} &\lesssim |I|^{\frac{q}{p}-1} \int_{S(I)} |\nabla f(x, t)|^q t^{q-1-q\alpha} dx dt \\ &\quad + \sum_{m=0}^{\infty} 2^{m(n-\frac{nq}{p}-\lambda)} |I_{m+1}|^{\frac{q}{p}-1} \int_{S(I_{m+1})} |\nabla f(x, t)|^q t^{q-1-q\alpha} dx dt \\ &\lesssim 1 + \sum_{m=0}^{\infty} 2^{m(n-\frac{nq}{p}-\lambda)} \lesssim 1. \end{aligned}$$

□

The following result is easily deduced from Theorem 3.4.

COROLLARY 3.5 *Let $0 \leq \alpha < \min(1, \frac{n}{q})$, $1 \leq q \leq p < \infty$ and $\lambda > n - \frac{nq}{p}$. The definitions of $H_{p,q}^{\alpha,\lambda}(\mathbb{R}_+^{n+1})$ are independent of the index λ .*

4. Harmonic function and Besov-Q spaces

In this section, by Theorems 2.8, 3.4 and 4.1, we extend the functions in $Q_{p,q}^\alpha(\mathbb{R}^n)$ to harmonic functions on \mathbb{R}_+^{n+1} . By Proposition 2.4, Theorems 3.4 and 4.2, we pull back the harmonic functions in $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$ to their relative trace function in $Q_{p,q}^\alpha(\mathbb{R}^n)$.

4.1. Poisson extension

In this subsection, we extend the functions in $Q_{p,q}^\alpha(\mathbb{R}^n)$ to harmonic functions in Carleson spaces. In fact,

THEOREM 4.1 Let $1 \leq q \leq p < \infty$ and $0 \leq \alpha < \min(1, \frac{n}{q})$. For any $f \in W_{p,q}^\alpha(\mathbb{R}^n)$, we have

$$f(x, t) =: P_t * f(x) \in C_{p,q}^\alpha(\mathbb{R}_+^{n+1}).$$

Proof By Theorem 3.4, it is enough to verify for $f \in W_{p,q}^\alpha(\mathbb{R}^n)$,

$$\sup_I |I|^{\frac{q}{p}-1} \int_{S(I)} |\nabla(P_t * f)(x)|^q t^{q-1-q\alpha} dx dt \lesssim \|f\|_{W_{p,q}^\alpha}.$$

For $i = 1, \dots, n$, define

$$C_{I,i} = |I|^{\frac{q}{p}-1} \int_{S(I)} \left| \frac{\partial f(x, t)}{\partial x_i} \right|^q t^{q-1-q\alpha} dx dt.$$

We only need to prove that

$$\sup_I C_{I,i} \lesssim \|f\|_{W_{p,q}^\alpha}^q, \quad i = 1, 2, \dots, n.$$

The kernel of $\frac{\partial f(x,t)}{\partial x_i}$ is $P_{i,t}(x)$. Let

$$\begin{cases} f_{\epsilon,j}(x) = \sum_{k \in \mathbb{Z}} a_{j,k}^\epsilon \Phi_{j,k}^\epsilon(x); \\ \frac{\partial f_{\epsilon,j}}{\partial x_i}(x, t) = P_{i,t} * f_{\epsilon,j}(x). \end{cases}$$

We obtain

$$\frac{\partial f(x, t)}{\partial x_i} = \sum_{\epsilon,j} \frac{\partial f_{\epsilon,j}(x, t)}{\partial x_i}.$$

By (2.5) and (2.6), we estimate $\partial f_{\epsilon,j}(x, t)/\partial x_i$ as follows. If $2^j t > 1$, then by integration by parts, we have

$$\frac{\partial f_{\epsilon,j}(x, t)}{\partial x_i} = \int_{\mathbb{R}^n} P_{i,\epsilon,t}(x-y) \sum_{k \in \mathbb{Z}} 2^{-j} a_{j,k}^\epsilon (I_\epsilon \Phi^\epsilon)_{j,k}(y) dy.$$

Hence, by (2.5), we can get

$$\begin{aligned} \left| \frac{\partial f_{\epsilon,j}(x, t)}{\partial x_i} \right| &\lesssim \int_{\mathbb{R}^n} |P_{i,t}(x-y)| \sum_{k \in \mathbb{Z}^n} |a_{j,k}^\epsilon| |\Phi_{j,k}^\epsilon(y)| dy \\ &\lesssim \sum_{k \in \mathbb{Z}^n} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+2)j} t (1 + |2^j x - k|)}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}}. \end{aligned}$$

If $2^j t \leq 1$, then

$$\frac{\partial f_{\epsilon,j}(x, t)}{\partial x_i} = \int_{\mathbb{R}^n} P_t(x-y) \sum_{k \in \mathbb{Z}^n} 2^j a_{j,k}^\epsilon \Phi_{j,k}^{\epsilon,i}(y) dy.$$

Hence, we have

$$\begin{aligned} \left| \frac{\partial f_{\epsilon, j}(x, t)}{\partial x_i} \right| &\lesssim \int_{\mathbb{R}^n} P_t(x-y) \sum_{k \in \mathbb{Z}^n} 2^j |a_{j, k}^\epsilon| |\Phi_{j, k}^{\epsilon, i}(y)| dy \\ &\lesssim \sum_{k \in \mathbb{Z}^n} |a_{j, k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1+|2^j x - k|^2)^{\frac{n+1}{2}}}. \end{aligned}$$

For $i = 1, \dots, n$ and $(\epsilon, j, k) \in \Lambda_n$, let

$$I(i, t, x, \epsilon, j, k) = \frac{\partial}{\partial x_i} \int P_t(x-y) \Phi_{j, k}^\epsilon(y) dy \tag{4.1}$$

be the function defined by (2.4). Then, we regroup the indices (ϵ, j, k) by I . Let $I_1 = 8I = \tilde{I}$ and $|I_1| = 2^{-nj}$. For $\tau \geq 1$, let I_τ be the cube which contains I_1 with $|I_\tau| = 2^{n\tau}|I_1|$. We divide the indices (ϵ, j, k) into three cases.

Case 1 $2^j t \geq 1$. For $l \in \mathbb{Z}^n$, define

$$\begin{cases} S_{-1, l} = \{(\epsilon, j, k) : 2^{-j}k \in 2^{-j}l + I_1, 2^j t \geq 1\}; \\ I_{-1, l}(i, t, x, I) = \sum_{(\epsilon, j, k) \in S_{-1, l}} |I(i, t, x, \epsilon, j, k)| |a_{j, k}^\epsilon|. \end{cases} \tag{4.2}$$

Case 2 $2^j t < 1 \leq 2^j \ell(I)$. For $l \in \mathbb{Z}^n$, define

$$\begin{cases} S_{0, l} = \{(\epsilon, j, k) : 2^{-j}k \in 2^{-j}l + I_1, 2^j t < 1 \leq 2^j \ell(I)\}; \\ I_{0, l}(i, t, x, I) = \sum_{(\epsilon, j, k) \in S_{0, l}} |I(i, t, x, \epsilon, j, k)| |a_{j, k}^\epsilon|. \end{cases} \tag{4.3}$$

Case 3 $2^j \ell(I) < 1$. For $l \in \mathbb{Z}^n$, define

$$\begin{cases} S_{\tau, l} = \{(\epsilon, j, k) : 2^{-j}k \in 2^{\tau-j}l + I_\tau, 2^j \ell(I) < 1\}; \\ I_{\tau, l}(i, t, x, I) = \sum_{(\epsilon, j, k) \in S_{\tau, l}} |I(i, t, x, \epsilon, j, k)| |a_{j, k}^\epsilon|. \end{cases} \tag{4.4}$$

Hence, we obtain that

$$\left| \frac{\partial f(x, t)}{\partial x_i} \right| \lesssim \sum_{\tau \geq -1, l \in \mathbb{Z}^n} I_{\tau, l}(i, t, x, I).$$

Now, we estimate the terms:

$$I_{i, \tau, l, I} = |I|^{\frac{q}{p}-1} \int_{S(I)} |I_{\tau, l}(i, t, x, I)|^q t^{q-1-q\alpha} dt.$$

We first estimate the case $\tau = -1$. At first, we assume $|l| \leq C$. We can see that

$$\begin{aligned} &|I|^{q/p-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in S_{-1, l}} |a_{j, k}^\epsilon| \frac{2^{(\frac{n}{2}+2)jt} (1+|2^j x - k|)}{(4jt^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \right\}^q t^{q-1-q\alpha} dx dt \\ &\lesssim |I|^{q/p-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in S_{-1, l}} |a_{j, k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j} (2^j t) (1+|2^j x - k|)}{(4jt^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \right\}^q t^{q-1-q\alpha} dx dt. \end{aligned}$$

Because $2^j t \geq 1$,

$$\frac{2^j t}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \lesssim \frac{1}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}}.$$

This gives

$$\begin{aligned} & \sum_{(\varepsilon, j, k) \in S_{-1, l}} |a_{j, k}^\varepsilon| \frac{2^{(\frac{n}{2}+1)j}}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \\ & \lesssim \sum_{j \geq -\log_2 t} 2^{(\frac{n}{2}+1)j} \left(\sum_{2^{-j} k \in 2^{-j} l + I_1} \frac{|a_{j, k}^\varepsilon|^q}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{1/q} \\ & \quad \times \left(\sum_{2^{-j} k \in 2^{-j} l + I_1} \frac{1}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{\frac{q-1}{q}} \\ & \lesssim \sum_{j \geq -\log_2 t} 2^{(\frac{n}{2}+1)j} (2^j t)^{-\frac{q-1}{q}} \left(\sum_{2^{-j} k \in 2^{-j} l + I_1} \frac{|a_{j, k}^\varepsilon|^q}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{1/q}. \end{aligned} \tag{4.5}$$

We obtain that

$$\begin{aligned} & |I|^{q/p-1} \int_{S(I)} \left\{ \sum_{(\varepsilon, j, k) \in S_{-1, l}} |a_{j, k}^\varepsilon| \frac{2^{(\frac{n}{2}+2)j} t (1 + |2^j x - k|)}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \right\}^q t^{q-1-q\alpha} dx dt \\ & \lesssim |I|^{q/p-1} \int_I \int_0^{\ell(I)} \sum_{\varepsilon, j \geq -\log_2 t} 2^{q(\frac{n}{2}+1)j} 2^{-j(q-1)} \\ & \quad \times \left(\sum_{k \in 2^{-j} l + 2^j I_1} \frac{|a_{j, k}^\varepsilon|^q (2^j t)^\delta}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right) t^{-q\alpha} dx dt. \end{aligned}$$

We change the order of summation and integration to get

$$\begin{aligned} & |I|^{q/p-1} \int_{S(I)} \left\{ \sum_{(\varepsilon, j, k) \in \Lambda_n} |a_{j, k}^\varepsilon| \frac{2^{(\frac{n}{2}+2)j} t (1 + |2^j x - k|)}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \right\}^q t^{q-1-q\alpha} dx dt \\ & \lesssim |I|^{q/p-1} \sum_{j \geq -\log_2 \ell(I)} 2^{q(\frac{n}{2}+1)j} 2^{-j(q-1)} 2^{j\delta} \sum_{\varepsilon, k} |a_{j, k}^\varepsilon|^q \\ & \quad \times \int_I \int_{2^{-j}}^{\ell(I)} \frac{1}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} t^{\delta-q\alpha} dx dt \\ & \lesssim |I|^{q/p-1} \sum_{j \geq -\log_2 \ell(I)} 2^{q(\frac{n}{2}+1)j} 2^{-j(q-1)} 2^{j\delta} 2^{-jn-j} 2^{-j(\delta-q\alpha)} \sum_{\varepsilon, k} |a_{j, k}^\varepsilon|^q \\ & \lesssim \|f\|_{W_{p, q}^\alpha}. \end{aligned}$$

If $|l| > C$, for $x \in I$, $|x - x_l| < \ell(I)$. On the other hand, $k \in 2^{j-j_l}l + 2^j I_1$ implies that $|k - 2^{j-j_l}l| \leq 2^j \ell(I_1) = 2^{j-j_l}$. We can see that $|2^j x - k| \sim 2^{j-j_l}(1 + |l|)$. Also for any fixed l , $\#\{k : k \in 2^{j-j_l}l + 2^j I_1\} = 2^{(j-j_l)n}$. From these estimates, we can deduce that

$$\sum_{k \in 2^{j-j_l}l + 2^j I_1} \frac{1}{(4j t^2 + |2^j x - k|^2)^{\frac{n+1}{2}}} \lesssim \frac{1}{2^j t (1 + |l|)^{n+1}},$$

where in the last inequality, we have used the facts that $8\ell(I) = 2^{-j_l}$ and $1 < 2^j t < 2^j \ell(I)$. Similar to the case of $|l| \leq C$, we still have

$$\begin{aligned} & |I|^{q/p-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in \Lambda_n} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+2)j} t (1 + |2^j x - k|)}{(4j t^2 + |2^j x - k|^2)^{\frac{n+3}{2}}} \right\}^q t^{q-1-q\alpha} dx dt \\ & \lesssim (1 + |l|)^{-q(n+1)} \|f\|_{W_{p,q}^\alpha}. \end{aligned}$$

We now estimate the case $\tau = 0$. We consider the case $|l| \leq C$.

$$I_{i,0,l,I} \lesssim |I|^{\frac{q}{p}-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q t^{q-1-q\alpha} dx dt.$$

Take $\delta > q - q\alpha > 0$. Applying Cauchy-Schwartz's inequality to k and j , respectively, we can obtain

$$\begin{aligned} & \left\{ \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q \\ & \lesssim \left\{ \sum_{\epsilon, -\log_2 \ell(I) \leq j < -\log_2 t} 2^{(\frac{n}{2}+1)j} \left(\sum_{k: (\epsilon, j, k) \in S_{0,l}} \frac{|a_{j,k}^\epsilon|^q}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{\frac{1}{q}} \right\}^q \\ & \lesssim \sum_{(\epsilon, j, k) \in S_{0,l}} 2^{q(\frac{n}{2}+1)j} \frac{|a_{j,k}^\epsilon|^q (2^j t)^{-\delta}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}}. \end{aligned}$$

The above estimate gives

$$\begin{aligned} I_{i,0,l,I} & \lesssim |I|^{\frac{q}{p}-1} \int_{S(I)} \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon|^q \frac{2^{(\frac{n}{2}+1)qj} (2^j t)^{-\delta}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} t^{q-1-q\alpha} dx dt \\ & \lesssim |I|^{\frac{q}{p}-1} \sum_{(\epsilon, j, k) \in S_{0,l}} 2^{qj(\alpha + \frac{n}{2}) - nj} (2^j \ell(I))^{q-q\alpha-\delta} |a_{j,k}^\epsilon|^q \\ & \lesssim \|f\|_{W_{p,q}^\alpha}, \end{aligned}$$

where in the last inequality, we have used the fact that $(\epsilon, j, k) \in S_{0,l}$ implies that $2^j \ell(I) \leq 1$.

We consider then the case $|l| > C$. At first, Cauchy-Schwartz's inequality gives

$$\left\{ \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q$$

$$\lesssim \left[\sum_{\epsilon, j < -\log_2 t} 2^{j(\frac{n}{2}+1)} \left(\sum_{k \in \mathbb{Z}^n} \frac{|a_{j,k}^\epsilon|^q}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{\frac{1}{q}} \left(\sum_{k \in \mathbb{Z}^n} \frac{1}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right)^{\frac{q-1}{q}} \right]^q.$$

Because $(\epsilon, j, k) \in S_{0,l}$, we can see that $|2^j x - k| \sim 2^{j-j_l} (1 + |l|)$ which implies that

$$\sum_{k \in \mathbb{Z}^n} \frac{1}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \sim (2^{j-j_l} (1 + |l|)^{n+1})^{-1}.$$

By the above estimate, we apply Cauchy-Schwartz's inequality to j and get

$$\left\{ \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q$$

$$\lesssim \sum_{(\epsilon, j, k) \in S_{0,l}} 2^{qj(\frac{n}{2}+1)} \frac{|a_{j,k}^\epsilon|^q}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \frac{(2^j t)^{-\delta}}{[2^{j-j_l} (1 + |l|)^{n+1}]^{q-1}}. \tag{4.6}$$

By (4.6), we can obtain

$$I_{i,0,l,t} \lesssim |l|^{\frac{q}{p}-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in S_{0,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q t^{q-1-q\alpha} dx dt$$

$$\lesssim |l|^{\frac{q}{p}-1} \int_{S(I)} \sum_{(\epsilon, j, k) \in S_{0,l}} \frac{|a_{j,k}^\epsilon|^q}{[2^{j-j_l} (1 + |l|)^{n+1}]^{q-1}} \frac{2^{(\frac{n}{2}+1)qj} (2^j t)^{-\delta}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} t^{q-1-q\alpha} dx dt$$

$$\lesssim (1 + |l|)^{-q(n+1)} |l|^{\frac{q}{p}-1} \sum_{(\epsilon, j, k) \in S_{0,l}} 2^{qj(\alpha+\frac{n}{2})-nj} |a_{j,k}^\epsilon|^q$$

$$\lesssim (1 + |l|)^{-q(n+1)} \|f\|_{W_{p,q}^\alpha}.$$

Thirdly, we estimate the case $\tau \geq 1$ where the number of (ϵ, j, k) is finite. We consider the case $|l| \leq C$.

$$I_{i,\tau,l,t} \lesssim |l|^{\frac{q}{p}-1} \int_{S(I)} \left\{ \sum_{(\epsilon, j, k) \in S_{\tau,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1 + |2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q t^{q-1-q\alpha} dx dt$$

$$\lesssim 2^{-q\delta\tau} |I_\tau|^{\frac{q}{p}-1} \sum_{(\epsilon, j, k) \in S_{\tau,l}} 2^{qj(\alpha+\frac{n}{2})-nj} |a_{j,k}^\epsilon|^q$$

$$\lesssim 2^{-q\delta\tau} \|f\|_{W_{p,q}^\alpha}.$$

We consider then the case $|l| > C$. Similar to the estimate (4.6), taking $0 < \delta < q - q\alpha$, we can get

$$\begin{aligned} I_{i,\tau,l} &\lesssim |I|^{\frac{q}{p}-1} \int_{S(I)} \left\{ \sum_{(\epsilon,j,k) \in S_{\tau,l}} |a_{j,k}^\epsilon| \frac{2^{(\frac{n}{2}+1)j}}{(1+|2^j x - k|^2)^{\frac{n+1}{2}}} \right\}^q t^{q-1-q\alpha} dx dt \\ &\lesssim |I|^{\frac{q}{p}-1} \int_{S(I)} \sum_{(\epsilon,j,k) \in S_{\tau,l}} \frac{|a_{j,k}^\epsilon|^q}{[2^{j-jl}(1+|l|)^{n+1}]^{q-1}} \frac{2^{(\frac{n}{2}+1)qj} (2^j t)^{-\delta}}{(1+|2^j x - k|^2)^{\frac{n+1}{2}}} t^{q-1-q\alpha} dx dt \\ &\lesssim 2^{-q\delta\tau} (1+|l|)^{-q(n+1)} |I_\tau|^{\frac{q}{p}-1} \sum_{(\epsilon,j,k) \in S_{\tau,l}} 2^{qj(\alpha+\frac{n}{2})-nj} |a_{j,k}^\epsilon|^q \\ &\lesssim 2^{-q\delta\tau} (1+|l|)^{-q(n+1)} \|f\|_{W_{p,q}^\alpha}, \end{aligned}$$

where we have used the fact that $(\epsilon, j, k) \in S_{\tau,l} \Leftrightarrow 2^{-j}k \in 2^{\tau-jl}l + I_\tau$ and $2^j \ell(I) < 1$. Take a positive number δ small enough. We repeat applying Cauchy-Schwartz's inequality

$$\begin{aligned} C_{I,i} &\lesssim \left\{ \sum_{\tau \geq -1, l \in \mathbb{Z}^n} 2^{-\tau\delta} (1+|l|)^{-(n+1)} \right\}^{q-1} \\ &\quad \times \sum_{\tau \geq -1, l \in \mathbb{Z}^n} 2^{(q-1)\tau\delta} (1+|l|)^{(q-1)(n+1)} |I|^{\frac{q}{p}-1} \int_{S(I)} |I_{\tau,l}(i, t, x, I)|^q t^{q-1-q\alpha} dt \\ &\lesssim \sum_{\tau \geq -1, l \in \mathbb{Z}^n} 2^{-\tau\delta} (1+|l|)^{-(n+1)} \|f\|_{W_{p,q}^\alpha}. \end{aligned}$$

□

4.2. Boundary value

The boundary value of a harmonic function in $C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$ may not be locally integrable. But we have

THEOREM 4.2 *Let $1 \leq q \leq p < \infty$ & $0 \leq \alpha < \min(1, \frac{n}{q})$. For any $f(x, t) \in C_{p,q}^\alpha(\mathbb{R}_+^{n+1})$, there exists a function $f \in W_{p,q}^\alpha(\mathbb{R}^n)$ such that*

$$f(x, t) = P_t * f(x).$$

Proof For simplicity, for any ϵ , let

$$(f, W_{p,q}^\alpha)_\epsilon(I) =: |I|^{\frac{q}{p}-1} \sum_{(j,k): I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} |f_{j,k}^\epsilon|^q.$$

We write

$$(f, W_{p,q}^\alpha)(I) = |I|^{\frac{q}{p}-1} \sum_{(\epsilon,j,k) \in \Lambda_n: I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} |f_{j,k}^\epsilon|^q \equiv \sum_\epsilon (f, W_{p,q}^\alpha)_\epsilon(I).$$

For $i = 1, \dots, n$ and any function f define

$$I_i f(x) = \int_{-\infty}^{x_i} f(x_1, \dots, x_{-1+i_\epsilon}, y, x_{1+i_\epsilon}, \dots, x_n) dy.$$

For $m \geq n + 8$, define $I_i^m f(x) = I_i I_i^{m-1} f(x)$. Let ϕ be a function in Lemma 2.2, we know $I_i^m \phi(x)$ is a C^{2n+8} function with compact support. For $\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_n) \in E_n$, denote by i_ϵ the smallest index such that $\epsilon_{i_\epsilon} = 1$. Let $\partial_\epsilon = \partial_{x_{i_\epsilon}}$ and $I_\epsilon \Phi^\epsilon(x) = I_{i_\epsilon} \Phi^\epsilon(x)$. Hence, we have

$$\begin{aligned} (f, W_{p,q}^\alpha)_\epsilon(I) &= |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} |f_{j,k}^\epsilon|^q \\ &= |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} \left| \left\langle \int_0^\infty \int_{\mathbb{R}^n} f(y,t) \phi_t(x-y) \frac{dt}{t} dy, \Phi_{j,k}^\epsilon \right\rangle \right|^q. \end{aligned}$$

We divide the integration on $(0, \infty)$ into two parts.

$$\begin{aligned} (f, W_{p,q}^\alpha)_\epsilon(I) &\lesssim |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} 2^{mqj} \\ &\quad \times \left| \left\langle \int_0^{2^{-j}} \int_{\mathbb{R}^n} \partial_1 f(y,t) (I_1^{m+1} \phi)_t(x-y) t^m dt dy, (\partial_1^m \Phi^\epsilon)_{j,k} \right\rangle \right|^q \\ &\quad + |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+\frac{n}{2})-nj} 2^{-(m+1)qj} \\ &\quad \times \left| \left\langle \int_{2^{-j}}^\infty \int_{\mathbb{R}^n} \partial_\epsilon f(y,t) (\partial_\epsilon^m \phi)_t(x-y) \frac{dt}{t^{m+1}} dy, (I_\epsilon^{m+1} \Phi^\epsilon)_{j,k} \right\rangle \right|^q \\ &=: J_0 + J_1, \end{aligned}$$

where

$$\begin{aligned} J_0 &= |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{mqj} \left\{ \int_0^{2^{-j}} \int_{\mathbb{R}^n \times \mathbb{R}^n} |\partial_1 f(y,t)| \left| (I_1^{m+1} \phi) \left(\frac{x-y}{t} \right) \right| \right. \\ &\quad \left. \times |(\partial_1^m \Phi^\epsilon)(2^j x - k)| t^{m-n} dt dx dy \right\}^q \end{aligned}$$

and

$$\begin{aligned} J_1 &= |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{-(m+1)qj} \left\{ \int_{2^{-j}}^\infty \int_{\mathbb{R}^n \times \mathbb{R}^n} |\partial_\epsilon f(y,t)| \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \right. \\ &\quad \left. \times \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| \frac{dt}{t^{m+n+1}} dx dy \right\}^q. \end{aligned}$$

We can see that

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^n \times \mathbb{R}^n} |\partial_1 f(y, t)| \left| \left(I_1^{m+1} \phi \right) \left(\frac{x-y}{t} \right) \right| \left| (\partial_1^m \Phi^\epsilon) (2^j x - k) \right| dx dy \right\}^q \\ & \lesssim 2^{-\frac{qnj}{q}} \int_{\mathbb{R}^n} \left\{ \int_{\mathbb{R}^n} |\partial_1 f(y, t)| \left| \left(I_1^{m+1} \phi \right) \left(\frac{x-y}{t} \right) \right| dy \right\}^q \left| (\partial_1^m \Phi^\epsilon) (2^j x - k) \right| dx \\ & \lesssim t^{n(q-1)} 2^{-n(q-1)j} \int_{\mathbb{R}^{2n}} |\partial_1 f(y, t)|^q \left| \left(I_1^{m+1} \phi \right) \left(\frac{x-y}{t} \right) \right| \left| (\partial_1^m \Phi^\epsilon) (2^j x - k) \right| dx dy. \end{aligned}$$

We first estimate the term J_0 . By Hölder's inequality, we can deduce that

$$\begin{aligned} J_0 & \lesssim |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(m+\alpha)-j(q-1)} \int_0^{2^{-j}} \int_{\mathbb{R}^{2n}} |\partial_1 f(y, t)|^q \left| \left(I_1^{m+1} \phi \right) \left(\frac{x-y}{t} \right) \right| \\ & \quad \times \left| (\partial_1^m \Phi^\epsilon) (2^j x - k) \right| t^{mq-n} dx dy dt. \end{aligned}$$

Notice that

$$u(y, t) = t^{-n} \sum_{I_{j,k} \subset I} \int_{\mathbb{R}^n} \left| \left(I_1^{m+1} \phi \right) \left(\frac{x-y}{t} \right) \right| \left| (\partial_1^m \Phi^\epsilon) (2^j x - k) \right| dx \in L^\infty(\tilde{I}).$$

Finally, we obtain

$$\begin{aligned} J_0 & \lesssim |I|^{\frac{q}{p}-1} \sum_{2^{nj}|I| \geq 1} 2^{qj(m+\alpha)-j(q-1)} \int_0^{2^{-j}} \int_{\tilde{I}} |\partial_1 f(y, t)|^q t^{mq} dy dt \\ & \lesssim |I|^{\frac{q}{p}-1} \int_{S(\tilde{I})} |\partial_1 f(y, t)|^q t^{q-1-q\alpha} dy dt. \end{aligned}$$

For sufficient small positive real number δ , we have

$$J_1 \lesssim \sum_{\tau=0}^{\infty} 2^{\delta\tau} J_{1,\tau},$$

where

$$\begin{aligned} J_{1,0} & = |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{-(m+1)qj} \left\{ \int_{2^{-j}}^{2^{\ell(I)}} \int_{\mathbb{R}^{2n}} \left| \partial_\epsilon f(y, t) \right| \right. \\ & \quad \left. \times \left| \left(\partial_\epsilon^m \phi \right) \left(\frac{x-y}{t} \right) \right| \left| \left(I_\epsilon^{m+1} \Phi^\epsilon \right) (2^j x - k) \right| \frac{dt}{t^{m+n+1}} dx dy \right\}^q \end{aligned}$$

and

$$\begin{aligned} J_{1,\tau} & = |I|^{\frac{q}{p}-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{-(m+1)qj} \left\{ \int_{2^{\tau-1}\ell(I)}^{2^\tau\ell(I)} \int_{\mathbb{R}^{2n}} \left| \partial_\epsilon f(y, t) \right| \right. \\ & \quad \left. \times \left| \left(\partial_\epsilon^m \phi \right) \left(\frac{x-y}{t} \right) \right| \left| \left(I_\epsilon^{m+1} \Phi^\epsilon \right) (2^j x - k) \right| \frac{dt}{t^{m+n+1}} dx dy \right\}^q. \end{aligned}$$

Via Hölder's inequality, a simple computation implies that

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^n \times \mathbb{R}^n} |\partial_\epsilon f(y, t)| \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx dy \right\}^q \\ & \lesssim t^{n(q-1)} 2^{-n(q-1)j} \int_{\mathbb{R}^{2n}} |\partial_\epsilon f(y, t)|^q \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx dy. \end{aligned}$$

Then, we can get

$$\begin{aligned} J_{1,0} & \lesssim |I|^{q/p-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{-(m+1)qj} 2^j \int_{2^{-j}}^{\ell(I)} \left\{ \int_{\mathbb{R}^{2n}} |\partial_\epsilon f(y, t)| \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \right. \\ & \quad \left. \times \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx dy \right\}^q \frac{dt}{t^{q(m+n)}} \\ & \lesssim |I|^{q/p-1} \sum_{I_{j,k} \subset I} 2^{qj\alpha} 2^{j-(m+1)qj} \int_{2^{-j}}^{\ell(I)} \int_{\mathbb{R}^{2n}} |\partial_\epsilon f(y, t)|^q \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \\ & \quad \times \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| t^{-mq-n} dx dy dt. \end{aligned}$$

Notice that

$$u(y, t) = t^{-n} \sum_{I_{j,k} \subset I} \int_{\mathbb{R}^n} \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx \in L^\infty(\tilde{I}).$$

We can obtain that

$$\begin{aligned} J_{1,0} & \lesssim |I|^{q/p-1} \sum_{2^{nj}|I| \geq 1} 2^{qj\alpha} 2^{j-(m+1)qj} \int_{2^{-j}}^{\ell(I)} \int_{\tilde{I}} |\partial_\epsilon f(y, t)|^q t^{-mq} dy dt \\ & \lesssim |I|^{q/p-1} \int_{S(\tilde{I})} |\partial_\epsilon f(y, t)|^q t^{q-1-q\alpha} dy dt \lesssim C. \end{aligned}$$

By a similar method, for $J_{1,\tau}$, we have

$$\begin{aligned} J_{1,\tau} & \lesssim |I|^{q/p-1} \sum_{I_{j,k} \subset I} 2^{qj(\alpha+n)-nj} 2^{-(m+1)qj} [2^\tau \ell(I)]^{-1} \int_{2^{\tau-1}\ell(I)}^{2^\tau \ell(I)} \left\{ \int_{\mathbb{R}^{2n}} |\partial_\epsilon f(y, t)| \right. \\ & \quad \left. \times \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx dy \right\}^q \frac{dt}{t^{q(m+n)}} \\ & \lesssim |I|^{q/p-1} \sum_{I_{j,k} \subset I} 2^{qj\alpha} 2^{-(m+1)qj} \{2^\tau l(I)\}^{-1} \int_{2^{\tau-1}\ell(I)}^{2^\tau \ell(I)} \int_{\mathbb{R}^{2n}} |\partial_\epsilon f(y, t)|^q \\ & \quad \times \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| t^{-mq-n} dx dy dt. \end{aligned}$$

We can see that

$$u(y, t) = t^{-n} \sum_{I_{j,k} \subset I} \int_{\mathbb{R}^n} \left| (\partial_\epsilon^m \phi) \left(\frac{x-y}{t} \right) \right| \left| (I_\epsilon^{m+1} \Phi^\epsilon) (2^j x - k) \right| dx \in L^\infty(\tilde{I}_\tau).$$

The above estimate gives

$$\begin{aligned}
 J_{1,\tau} &\lesssim |I|^{\frac{q}{p}-1} \sum_{2^{nj}|I|\geq 1} 2^{qj\alpha} 2^{-(m+1)qj} [2^\tau \ell(I)]^{-1} \int_{2^{\tau-1}\ell(I)}^{2^\tau \ell(I)} \int_{\tilde{I}_\tau} |\partial_\epsilon f(y, t)|^q t^{-mq} dy dt \\
 &\lesssim 2^{n\tau(1-\frac{q}{p})+\tau(q\alpha-mq-q)} |I_\tau|^{\frac{q}{p}-1} \int_{S(\tilde{I}_\tau)} |\partial_\epsilon f(y, t)|^q t^{q-1-q\alpha} dy dt \\
 &\lesssim 2^{n\tau(1-\frac{q}{p})+\tau(q\alpha-mq-q)}.
 \end{aligned}$$

□

Funding

Pengtao Li's research is supported by NSFC [grant number 11171203], [grant number 11201280]; New Teacher's Fund for Doctor Stations, Ministry of Education [grant number 20114402120003]; Foundation for Distinguished Young Talents in Higher Education of Guangdong, China, LYM11063. Qixiang Yang's research is supported in part by NSFC [grant number 11271209]. Tao Qian's research is supported in part by MYRG116(Y1-L3)-FST13- QT; MYRG115(Y1-L4)-FST13-QT; and FDCT 098/2012/A3.

References

- [1] Fabes E, Johnson R, Neri U. Spaces of harmonic functions representable by Poisson integrals of functions in BMO and $\mathcal{L}_{p,\lambda}$. *Indiana Univ. Math. J.* 1976;25:159–170.
- [2] Peetre J. *New thoughts on Besov spaces*. Durham: Duke Univ. Math. Ser., Duke Univ. Press; 1976.
- [3] John F, Nirenberg L. On functions of bounded mean oscillation. *Comm. Pure Appl. Math.* 1961;14:415–426.
- [4] Essen M, Janson S, Peng L, Xiao J. Q spaces of several real variables. *Indiana Univ. Math. J.* 2000;49:575–615.
- [5] Wu Z, Xie C. Decomposition theorems for Q_p spaces. *Ark. Mat.* 2002;40:383–401.
- [6] Meyer Y. *Wavelets and operators*. Vol. 37, Cambridge studies in advanced mathematics. Cambridge: Cambridge University Press; 1992.
- [7] Meyer Y. La minimalité de l'espace de Besov et la continuité de l'opérateurs par l'intégrales singulieres, monografias de Matematicas [The minimality of Besov space and the continuity of the singular integral operators]. *Universidad autonoma de madrid.* 1986;4:1–37.
- [8] Xiao J. Homothetic variant of fractional Sobolev space with application to Navier-Stokes system. *Dyn. Partial Differ. Equ.* 2007;2:227–245.
- [9] Cui L, Yang Q. On the generalized Morrey spaces. *Siberian Math. J.* 2005;46:133–141.
- [10] Lin C, Yang Q. Semigroup characterization of Besov type Morrey spaces and well-posedness of generalized Navier-Stokes equations. *J. Differ. Equ.* 2013;254:804–846.
- [11] Yang D, Yuan W. A new class of function spaces connecting Triebel-Lizorkin spaces and Q spaces. *J. Funct. Anal.* 2008;255:2760–2809.
- [12] Wojtaszczyk P. *A mathematical introduction to wavelets*. Vol. 37, London mathematical society student texts. New York: Cambridge University Press; 1997.
- [13] Yang Q. *Wavelet and distribution*. Beijing: Beijing Science and Technology Press; 2002.
- [14] Dafni G, Xiao J. Some new tent spaces and duality theorem for fractional Carleson measures and $Q_\alpha(\mathbb{R}^n)$. *J. Funct. Anal.* 2004;208:377–422.

- [15] Dafni G, Xiao J. The dyadic structure and atomic decomposition of Q spaces in several real variables. *Tohoku Math. J.* 2005;57:119–145.
- [16] Li P, Yang Q. Well-posedness of Quasi-geostrophic equations with data in Besov- Q spaces. *Nonlinear Anal.* 2014;94:243–258.
- [17] Li P, Yang Q. Wavelets and the well-posedness of incompressible magneto-hydrodynamic equations in Besov type Q -space. *J. Math. Anal. Appl.* 2013;405:661–686.
- [18] Zhao R. On a general family of function spaces. *Ann. Acad. Sci. Fenn. Math. Diss.* 1996;105. 56 p.
- [19] Chaisuriya P, Reséndis O, Lino F, Tovar S, Luis M, Zhao R. A new class of analytic function with measures. *Complex Var. Elliptic Equ.* 2013;58:1145–1160.
- [20] Lindström M, Palmberg N. Duality of a large family of analytic function spaces. *Ann. Acad. Sci. Fenn. Math.* 2007;32:251–267.
- [21] Pérez-González F, Rättyä J. Forelli-Rudin estimates, Carleson measures and $F(p, q, s)$ -functions. *J. Math. Anal. Appl.* 2006;315:394–414.
- [22] Rättyä J. n th derivative characterisations, mean growth of derivatives and $F(p, q, s)$ Bull. Aust. Math. Soc. 2003;68:405–421.
- [23] Zhu K. A class of Möbius invariant function spaces. *Illinois J. Math.* 2007;51:977–1002.
- [24] Wang Y, Xiao J. Homogeneous Campanato-Sobolev classes. *Appl. Comput. Harmon. Anal.* 2014. doi:10.1016/j.acha.2014.09.002.
- [25] Jiang R, Xiao J, Yang D. Towards spaces of harmonic functions with traces in square Campanato space and its scaling invariant. 2013. arXiv:1309.7576.